

CCHMC Epic Order Set

▼ Nursing

▼ Nursing


Appropriate NGT tube size by age:

	Wt (kg)	NGT Size (Fr)	Suggested
Newborn/ Infant	3 - 9	5 - 8	8
Toddler	10 - 11	8 - 10	10
Small Child	12 - 14	10	12
Child	15 - 18	10 - 12	14
Child	19 - 23	12 - 14	16
Large Child/ Adult	>24	12 - 18	18

Insert NG Tube

Routine, ONCE, today at 0743, For 1 occurrence

NG to Suction

 Routine, ONCE, today at 0743, For 1 occurrence

Clamp NGT after gastrografin administration

Routine, ONCE, today at 0743, For 1 occurrence

Contact surgery team if patient has nausea, vomiting, abdominal pain while NGT is clamped after gastrografin administration

Routine

▼ Medications

▼ Patients < 8 years of age

diatrizoate meglumine & sodium (GASTROGRAFIN) 66-10 % solution

50 mL, NG tube, AS DIRECTED, see PRN comment, Nurse will add 50 mL of water to gastrografin for NGT administration by surgery team for 100 mL total volume, for 1 dose, Nurse will add 50 mL of water to gastrografin for NGT administration by surgery team for 100 mL total volume

▼ Patients >= 8 years of age


diatrizoate meglumine & sodium (GASTROGRAFIN) 66-10 % solution

100 mL, NG tube, AS DIRECTED, see PRN comment, Nurse will add 100 mL of water to gastrografin for NGT administration by surgery team for 200 mL total volume, for 1 dose, Nurse will add 100 mL of water to gastrografin for NGT administration by surgery team for 200 mL total volume


▼ Radiology

▼ Radiology

RAD Abdomen 1V

 Routine, ONCE, today at 0743, For 1 occurrence, to confirm NGT in stomach

RAD Abdomen 1V

 Routine, ONCE, today at 1743, For 1 occurrence, nonportable at 10 hours post Gastrografin administration

▼ Ad hoc Orders

 Search