SmartSet: PEDIATRIC SURGERY - APPENDICITIS PRE-OP (ID:692)

General Information

Display name: PEDIATRIC SURGERY - APPENDICITIS PRE-OP

Type: General

Merge priority: 0

Version comment:

Content source:

Synonyms: 1. PED SURG

SmartSet notes:

Description:

Web information:

	Title	URL
1.	Appendicitis Algorithm	http://portal2/sites/coex/clinical-qlty-excellence/clinical-qlty-effectiveness/Clinical%20Care%20Guidelines%20in%20Practice/Appendicitis/Appendicitis%20Algorithm%20FINAL.pdf
2.	Appendicitis Handout	http://portal2/sites/coex/clinical-qlty-excellence/clinical-qlty-effectiveness/Clinical%20Care%20Guidelines%20in%20Practice/Appendicitis/Appendicitis%20-%20Brochure.pdf

Questionnaire:

Configuration

Admission Orders

Admission Orders

☐ ADMIT TO INPATIENT	Admit To: Effective Date: 6/3/2022 Effective Time: Change to Attending Provider: Discharge Time: Discharge at 12:00 AM
☑ ADMIT TO OBSERVATION	Admit To: Effective Date: 6/3/2022 Effective Time: Change to Attending Provider: Discharge Time: Discharge at 12:00 AM, Observation

Routine Orders

Routine

☑ VITAL SIGNS	Routine, EVERY 4 HOURS
☑ ACTIVITY ORDER	Routine, AD LIB
☑ STRICT INTAKE AND OUTPUT	Routine, EVERY 4 HOURS
☑ CARDIAC/RESPIRATORY MONITORING IP UNITS	Routine
☑ PULSE OX CONTINUOUS IP-NURSING	Routine

OR CASE REQUEST ORDER

☐ SURGICAL/PROCEDURAL ADD-ON CASE REQUEST	Laparoscopic appendectomy

Diet Orders

Diet

☑ NPO (Enter appropriate starting date and time in NPO Order)	CONTINUOUS, Routine NPO Except: Sips with Meds

Nursing Orders

Nursing

☑ Patient is on Appendicitis Clinical Care Guideline	Routine, CONTINUOUS For 1
☑ NURSING COMMUNICATION	Please prepare patient for possible discharge the day of surgery (family to arrange transportation in advance).

☑ PROVIDE PATIENT WITH APPENDICITIS BROCHURE - (FOR UNCOMPLICATED APPENDICITIS PATIENTS ONLY) — Details

☑ NOTIFY MD/PROVIDER — Routine, AS NEEDED, Notify Pediatric Surgery MD/provider for: Temperature >= 38.5, O2 Sat < 92%, UOP < 1 mg/kg/hr, persistent pain, or with any questions or concerns.

☑ NURSING COMMUNICATION - MyChart Activation — Please ask caregiver if the patient's MyChart account has been activated. If not activated, please follow the steps to activate the account. Also, it's important to have them download the MyChart app onto their mobile phone, confirm notifications are turned on and click on the patient's name once logged in.

Medication Orders

IV Fluid

☑ dextrose 5% and 0.9% NACL with 20 mEq/L potassium chloride solution — Intravenous, CONTINUOUS

IV Flush

☑ sodium chloride 0.9 % flush syringe — Intravenous, AS NEEDED, Medication Flush

Acetaminophen

☐ acetaminophen (TYLENOL) suspension — 15 mg/kg/dose, Oral, EVERY 6 HOURS
☐ acetaminophen (TYLENOL) tablet — 15 mg/kg/dose, Oral, EVERY 6 HOURS

Morphine - for breakthrough pain

☐ morphine (DURAMORPH) injection — 0.05 mg/kg/dose, Intravenous, for 5 Minutes, EVERY 2 HOURS PRN, Severe Pain

Nausea/Vomiting

☐ ondansetron (ZOFRAN) injection — 0.1 mg/kg/dose, Intravenous, for 5 Minutes, EVERY 6 HOURS PRN, Nausea, Vomiting

Antibiotics

Antibiotics

☐ ceftriaxone (ROCEPHIN) syringe pump — 50 mg/kg/dose, Intravenous, for 15 Minutes, ONCE, For 1 Doses, For 1 Doses
☐ metronidazole (FLAGYL) injection — 30 mg/kg/dose, Intravenous, for 1 Hours, ONCE, For 1 Doses, For 1 Doses

Antibiotics - PCN Allergic

| ☐ ciprofloxacin (CIPRO) injection | 15 mg/kg/dose, Intravenous, for 1 Hours, ONCE, For 1 Doses, For 1 Doses |
| ☐ metronidazole (FLAGYL) injection | 30 mg/kg/dose, Intravenous, for 1 Hours, ONCE, For 1 Doses, For 1 Doses |

Consult Orders

Consults

| ☐ CHILD LIFE CONSULT | Routine, ONCE, Starting S+00 |
| ☐ SOCIAL WORK CONSULT | Routine, ONCE, Starting S+00 |

Criteria

Suggestions:

Filter:

Restrict SmartSet: ☐

Settings

Discontinue action:

Deselect sections for Pended/Held orders: ☐

Pended/Held orders display:

Release date: Use System Definitions Setting

Disallow user override: ☑